

Partition theorems for Ketonen-Solovay largeness

Quentin Le Houérou

Ludovic Patey

February 8, 2026

Abstract

We develop the framework of α -largeness introduced by Ketonen and Solovay, by proving a partition theorem for α -large sets with $\alpha < \epsilon_0$ which generalizes theorems from Ketonen and Solovay and from Bigorajska and Kotlarski. We also prove that for every ω^{nk+3} -large set X with $\min X \geq 18$, every coloring $f : [X]^2 \rightarrow k$ admits an ω^n -large f -homogeneous subset. This bound is tight, up to an additive constant.

1 Introduction

The celebrated Paris-Harrington theorem states that Peano arithmetic does not prove the existence of ω -large homogeneous sets for arbitrary instances of Ramsey's theorem. Here, a finite set X is ω -large if $\text{card } X > \min X$. Ketonen and Solovay [12] defined a notion of α -largeness for $\alpha < \epsilon_0$ to quantify the size of finite sets over weak theories, and related it to the Wainer hierarchy of fast-growing functions [27]. More precisely, any fundamental sequence system $(\{\alpha\}(n))_{n \in \mathbb{N}}$ for the ordinals $\alpha < \epsilon_0$ induces a notion of largeness defined as follows: a finite set of integers $\{x_0 < \dots < x_{k-1}\}$ is α -large if, letting $\alpha_0 = \alpha$ and $\alpha_{s+1} = \{\alpha_s\}(x_s)$, the ordinal α_{k-1} equals 0.

Since then, variants of Ketonen and Solovay's notion of largeness have been extensively studied, both from a combinatorial [3, 1, 16, 2, 5] and a proof-theoretic perspective [24, 15, 22, 13, 23, 26], to prove partial conservation theorems over subsystems of second-order arithmetic for $\forall\Sigma_2^0$ -sentences and non-speedup theorems, where a $\forall\Sigma_n^0$ -formula is of the form $\forall X\varphi(X)$ where $\varphi(X)$ is Σ_n^0 . More recently, parameterized versions of largeness have been defined to prove partial conservation theorems over $\text{RCA}_0 + \text{B}\Sigma_2^0$ for $\forall\Sigma_3^0$ -sentences [17, 18].

The framework of α -largeness is however relatively scattered: Ketonen and Solovay [12] first proved some general structural theorems about α -largeness for $\alpha < \epsilon_0$. Then Bigorajska and Kotlarski [2, 15, 3, 1] published a series of articles on a variant of α -largeness, say α -largeness*, introduced by Ratajczyk [24] based on the Hardy hierarchy of fast-growing functions [10]. They proved in particular

a general pigeonhole theorem for α -largeness* with $\alpha < \epsilon_0$. A framework for proving lower bounds for α -largeness* was developed by multiple authors [3, 16]. Both notions of largeness are related (in particular, any α -large set is α -large* for $\alpha \geq \omega$), but the translation of structural theorems from one notion to the other yields sub-optimal statements. Independently, the study of partial conservation theorems over $\mathbb{I}\Sigma_1$ motivated the development of the framework of α -largeness for $\alpha < \omega^\omega$ [22, 13, 23, 26]. In particular, Kołodziejczyk and Yokoyama [14] proved that for every $\omega^{144(n+1)}$ -large set F and every 2-coloring $f : [F]^2 \rightarrow 2$, there exists an ω^n -large f -homogeneous set. For this, they proved multiple structural theorems in the restricted setting of $\alpha < \omega^\omega$.

In this article, we pursue the general study of Ketonen and Solovay's notion of largeness, by proving the following general partition theorem, where \oplus denotes the natural (or Hessenberg) sum over ordinals, and a set F is *at most α -large* if $F \setminus \{\max F\}$ is α -small.

Main Theorem 1.1. *Let $\beta, \gamma < \epsilon_0$ be two ordinals, and $B, C \subseteq \mathbb{N}$ be at most β -large and γ -large respectively. Then $B \cup C$ is at most $(\beta \oplus \gamma)$ -large.*

This theorem has several consequences, including a generalization of Ketonen and Solovay [12, Lemma 4.6] and pigeonhole generalizations of Bigorajska and Kotlarski [1, Theorem 12] for α -largeness. For this, we introduce a Hardy-like hierarchy of fast-growing functions, and adapt and generalize the techniques of Bigorajska and Kotlarski.

Our second contribution is a tight upper bound (up to additive constant) of the closure of α -largeness under Ramsey's theorem for pairs, for $\alpha < \omega^\omega$. We say that a set F is RT_k^2 - α -large if for every k -coloring of the pairs $[F]^2$, there is an α -large f -homogeneous subset. We prove the following theorem:

Main Theorem 1.2. *Let $n, k \geq 1$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{kn+3} -large and $\min X \geq 17$, then it is RT_k^2 - ω^n -large.*

Note that in the case $k = 2$, we obtain ω^{2n+3} -largeness, improving over the $\omega^{144(n+1)}$ -largeness upper bound of Kołodziejczyk and Yokoyama [14]. This bound is tight, in the sense that Kotlarski et al. [16, Theorem 5.4] proved that if a set is RT_k^2 - ω^n -large*, then it is ω^{kn} -large*. Translating in the α -largeness framework, this shows that every RT_k^2 - ω^n -large is ω^{kn-1} -large.

The proof of Main Theorem 1.2 goes through the computation of upper bounds of closure for two intermediate combinatorial theorems, namely, the restriction of Ramsey's theorem for pairs to transitive colorings (trRT_k^2) and a generalization of the Erdős-Moser theorem to k -colorings (fEM) due to Towsner and Yokoyama [26]. We prove in particular that trRT_k^2 is solely responsible for the lower bound of RT_k^2 - ω^n -largeness, in that every ω^{n+3} -large set X with $\min X \geq 7$ is fEM - ω^n -large (Corollary 3.15). This later result improves the ω^{18n} -largeness upper bound of Towsner and Yokoyama for fEM - ω^n -largeness.

The remainder of this article is divided into two parts, as follows: In Section 2, we define and study Ketonen and Solovay's notion of α -largeness for $\alpha < \epsilon_0$ and prove Main Theorem 1.1 and its consequences. Then, in Section 3, we restrict ourselves to α -largeness for $\alpha < \epsilon_0$ and study the closure of

α -largeness under multiple combinatorial statements, including Ramsey's theorem for pairs. The section culminates with the proof of Main Theorem 1.2.

2 Largeness below ϵ_0

Every ordinal $\alpha < \epsilon_0$ admits a unique *Cantor normal form*

$$\alpha = \omega^{\alpha_s} \cdot a_s + \cdots + \omega^{\alpha_0} \cdot a_0$$

for some $\alpha_0 < \alpha_1 < \cdots < \alpha_s < \alpha$ and $a_0, \dots, a_s \in \mathbb{N} \setminus \{0\}$. Given two ordinals $\alpha, \beta < \epsilon_0$, we write $\beta \gg \alpha$ if either α or β equals 0, or if the smallest exponent of β is at least the largest exponent of α in their Cantor normal form. If $\alpha \neq 0$, we shall also use its *short Cantor normal form* $\alpha = \beta + \omega^\gamma$ with $\beta \gg \omega^\gamma$.

Definition 2.1. *Given a non-zero ordinal $\alpha < \epsilon_0$ with short Cantor normal form $\alpha = \beta + \omega^\gamma$, and $x \in \mathbb{N}$, let*

$$\{\alpha\}(x) = \begin{cases} \beta & \text{if } \gamma = 0, \\ \beta + \omega^{\gamma-1} \cdot x & \text{if } \gamma \text{ is successor,} \\ \beta + \omega^{\{\gamma\}(x)} & \text{if } \gamma \text{ is limit.} \end{cases}$$

Finally, let $\{0\}(x) = 0$.

The \gg relation admits two desirable properties. First, if $\beta \gg \alpha$, then the sum $\alpha + \beta$ coincides with the natural sum (or Hessenberg sum) $\alpha \oplus \beta$ corresponding to the component-wise sum of the multiplicative factors. Second, if $\beta \gg \alpha$ and $\alpha \neq 0$, then the operation $\{\beta + \alpha\}(x)$ can be reduced to a computation of $\{\alpha\}(x)$:

Lemma 2.2 (Ketonen and Solovay [12]). *Let $\alpha, \beta < \epsilon_0$ be such that $\beta \gg \alpha$ and $\alpha \neq 0$. Then $\{\beta + \alpha\}(x) = \beta + \{\alpha\}(x)$.*

In general, fundamental sequences are not compatible with ordinal inequality, in the sense that there exist some ordinals $\beta > \alpha$ and some $x \in \mathbb{N}$ such that $\{\beta\}(x) < \{\alpha\}(x)$. For instance, if $\beta = \omega$ and $\alpha = 10$, then $\{\omega\}(1) = 1 < \{10\}(1) = 9$. However, this is the case if x is larger than every coefficient appearing in the Cantor normal form of α . This motivates the definition of the pseudo-norm:

Definition 2.3. *The pseudo-norm $\text{psn}(\alpha)$ of α of cantor normal form $\omega^{\alpha_s} \cdot a_s + \cdots + \omega^{\alpha_0} \cdot a_0$ is defined as*

$$\text{psn}(\alpha) = \max(\text{psn}(\alpha_0), \dots, \text{psn}(\alpha_s), a_0, \dots, a_s)$$

The following lemma is an immediate consequence of Bigorajska and Kotlarski [3, Lemma 2.1].

Lemma 2.4 (Bigorajska and Kotlarski [3]). *Let $\alpha < \beta < \epsilon_0$ and $x \in \mathbb{N}$ such that $\text{psn}(\alpha) < x$. Then $\{\beta\}(x) \geq \alpha$ (with equality if only if $\beta = \alpha + 1$).*

The following easy lemma relates $\text{psn}(\{\alpha\}(n))$ to $\text{psn}(\alpha)$ for any $n \in \mathbb{N}$.

Lemma 2.5. *Let $\alpha < \epsilon_0$ and let $n \in \mathbb{N}$. Then $\text{psn}(\alpha) - 1 \leq \text{psn}(\{\alpha\}(n)) \leq \max\{\text{psn}(\alpha), n\}$*

Proof. The result is clear for $\alpha = 0$, then, if $\alpha \neq 0$, the operation $\{\alpha\}(n)$ can only add n 's and decrease coefficients by 1 in the (recursive) Cantor normal form of α . □

The notion of fundamental sequence generalizes to finite sets as follows:

Definition 2.6 (Ketonen and Solovay). *Given an ordinal $\alpha < \epsilon_0$ and a finite set $X = \{x_0 < \dots < x_s\}$, let*

$$\{\alpha\}(X) = \{\dots \{\{\alpha\}(x_0)\}(x_1) \dots\}(x_s)$$

A set X is α -large if $\{\alpha\}(X) = 0$, otherwise, it is α -small.

Accordingly, a finite set X is *exactly α -large* if it is α -large and $X \setminus \{\max X\}$ is α -small. A finite set X is *at most α -large* if $X \setminus \{\max X\}$ is α -small. By convention, we consider that \emptyset is at most 0-large.

Example 2.7. *Any finite set $X \subseteq \mathbb{N}$ is $\text{card } X$ -large. A finite set X is ω -large if $\text{card } X > \min X$. A finite set X is $\omega^\beta \cdot n$ -large iff X contains n ω^β -large subsets $X_0 < \dots < X_{n-1} \subseteq X$, where $X < Y$ means $\forall x \in X \forall y \in Y x < y$.*

Note that a set X might be α -large, $\alpha + 1$ -small, but not at most α -large. For instance, $\{3, 5, 6, 7, 8\}$ is ω -large, not $\omega + 1$ -large, and not at most ω -large.

2.1 Hardy-like hierarchy

Ratajczyk [24, Section 5] defined a notion of largeness based on the Hardy hierarchy of functions [10]. As noted by Ratajczyk, this notion is slightly different from Ketonen and Solovay's α -largeness, and was extensively studied by Bigorajska and Kotlarski [1, 3, 15]. We now define a Hardy-like hierarchy whose induced notion of largeness corresponds exactly to Ketonen and Solovay's definition.

Definition 2.8. *Fix an increasing function $h : A \rightarrow \mathbb{N}$ for some $A \subseteq \mathbb{N}$ with $h(a) > a$ for every $a \in A$. We define inductively a sequence of partial functions h_α for every $\alpha < \epsilon_0$ as follows: for $x \in \mathbb{N}$ and $\alpha > 0$, let $h_0(x) = x$ and let $h_\alpha(x) = h_{\{\alpha\}(x)}(h(x))$ if $h(x) \in \text{dom}(h_{\{\alpha\}(x)})$ and $x \in \text{dom}(h)$, otherwise let $h_\alpha(x)$ undefined. We write $h_\alpha(x) \downarrow$ is $x \in \text{dom } h_\alpha$ and $h_\alpha(x) \uparrow$ otherwise.*

Note that for every $\alpha < \epsilon_0$ and $x \in \text{dom } h_\alpha$, there is some $n \in \mathbb{N}$ such that $h_\alpha(x) = h^{(n)}(x)$. However, this n depends on x : if $\alpha \geq \omega$, there is no n such that $h_\alpha = h^{(n)}$.

Remark 2.9. In the definition of the original Hardy hierarchy [10] based on h , if α is limit, then $h_\alpha(x) = h_{\{\alpha\}(x)}(x)$, while in our Hardy-like hierarchy, we define $h_\alpha(x) = h_{\{\alpha\}(x)}(h(x))$. This variation reflects the main difference between the notion of largeness from Ratajczyk [24, Section 5] and the one of Ketonen and Solovay [12]. In the former case, a set A is α -large for a limit ordinal α if it is $\{\alpha\}(\min A)$ -large, while in the latter case, a set A is α -large if $A \setminus \{\min A\}$ is $\{\alpha\}(\min A)$ -large. Some further differences exist: in the finite case, a set A is n -large in the sense of Ratajczyk if $\text{card } A \geq n + 1$ while in Ketonen and Solovay's sense, a set A is n -large iff $\text{card } A \geq n$. As a consequence, under both definitions, a set A is ω -large iff $\text{card } A > \min A$.

Lemma 2.10. If $h_\alpha(x) \downarrow$ for some $\alpha < \epsilon_0$ and $x \in \mathbb{N}$, then $h_\alpha(x) \geq \text{psn}(\alpha) + x$.

Proof. By induction on α . For $\alpha = 0$, let $x \in \mathbb{N}$ be such that $h_0(x) \downarrow$, then $h_0(x) = x = \text{psn}(0) + x$. Let $\alpha < \epsilon_0$ and assume the property to hold for every $\alpha' < \alpha$. Let $x \in \mathbb{N}$ be such that $h_\alpha(x) \downarrow$, then $h_\alpha(x) = h_{\{\alpha\}(x)}(h(x))$. By the inductive hypothesis, we have $h_{\{\alpha\}(x)}(h(x)) \geq \text{psn}(\{\alpha\}(x)) + h(x)$, but, as $h(x) \geq x + 1$ and as $\text{psn}(\{\alpha\}(x)) \geq \text{psn}(\alpha) - 1$ by Lemma 2.5, we have $h_{\{\alpha\}(x)}(h(x)) \geq \text{psn}(\alpha) + x$. \square

Fix an increasing function $h : A \rightarrow \mathbb{N}$ for some $A \subseteq \mathbb{N}$, with $h(x) > x$ for every $x \in A$. The Hardy-like iterations of h yield faster-growing functions on a restricted domain, that is, if $\alpha > 0$, then $\text{dom } h_\alpha \subseteq \text{dom } h$ and h_α dominates h on $\text{dom } h_\alpha$ (see Lemma 2.14(4)).

On the other hand, given two ordinals $\alpha \leq \beta < \epsilon_0$ and some $x \in \text{dom } h_\beta$, it is not true in general that $x \in \text{dom } h_\alpha$, nor that if $x \in \text{dom } h_\beta \cap \text{dom } h_\alpha$, then $h_\beta(x) \geq h_\alpha(x)$. However, if x is sufficiently large – $x > \text{psn}(\alpha)$ suffices –, then this is the case (see Lemmas 2.12 and 2.14). However, the $x > \text{psn}(\alpha)$ hypothesis is too strong for our purpose, so we will use a more general relation $\beta \Rightarrow_x \alpha$ which depends on both β and α , and such that if $\beta \Rightarrow_x \alpha$, then $\beta \geq \alpha$ and $h_\beta(x) \geq h_\alpha(x)$.

Definition 2.11 (Ketonen and Solovay). *Given two ordinals $\alpha, \beta < \epsilon_0$ and $x \geq 0$, let $\beta \Rightarrow_x \alpha$ if either $\beta = \alpha$, or $\{\beta\}(x) \Rightarrow_x \alpha$.*

Bigorajska and Kotlarski [1, Lemma 2(viii)] proved that the definition is unchanged if one replaces $\{\beta\}(x)$ by $\{\beta\}(y)$ for some $y \leq x$. As one expects from the previous discussion, the relation $\beta \Rightarrow_x \alpha$ coincides with the inequality $\beta \geq \alpha$ whenever $x > \text{psn}(\alpha)$. The whole interest of this relation is therefore in the case $x \leq \text{psn}(\alpha)$.

The following lemma strengthens Ketonen and Solovay [12, Proposition 2.8.1] and is an immediate consequence of Bigorajska and Kotlarski [3, Lemma 2.1]. It appears in this form in the posthumous book of Kotlarski [15, Lemma 1.4.3].

Lemma 2.12 (Kotlarski [15]). *Let $\alpha \leq \beta < \epsilon_0$ and $x \in \mathbb{N}$ such that $\text{psn}(\alpha) < x$, then $\beta \Rightarrow_x \alpha$.*

Ketonen and Solovay [12] proved many properties about the relation \Rightarrow_x . The following lemma compiles a list of basic facts that shall be useful to prove our main theorem.

Lemma 2.13 (Ketonen and Solovay [12]). *Let $\alpha, \beta, \lambda < \epsilon_0$ and $x \geq 0$.*

- (1) *If $\lambda \gg \alpha$ and $\alpha \Rightarrow_x \beta$, then $(\lambda + \alpha) \Rightarrow_x (\lambda + \beta)$;*
- (2) *If $k < \ell \in \mathbb{N}$, then $\omega^\alpha \cdot \ell \Rightarrow_x \omega^\alpha \cdot k$;*
- (3) *If $\alpha \Rightarrow_x \beta$ and $\alpha > \beta + 1$, then $\alpha \Rightarrow_{x+1} \beta + 1$.*
- (4) *If $\alpha \Rightarrow_x \beta$ and $x \geq 1$, then $\omega^\alpha \Rightarrow_x \omega^\beta$;*
- (5) *If $x \geq 1$ and $k < \ell \in \mathbb{N}$, then $\{\alpha\}(\ell) \Rightarrow_x \{\alpha\}(k)$.*

Proof. (1) is [12, Lemma 1, part 2.4], (2) is [12, Lemma 3, part 2.4], (3) is [12, Lemma 2, part 2.6], (4) is [12, Lemma 5, part 2.4], (5) is [12, Theorem 2.4] for α limit. If α is 0 or successor, then $\{\alpha\}(\ell) = \{\alpha\}(k)$. \square

The following lemma states that the \Rightarrow_x relation is a sufficient hypothesis for the Hardy-like hierarchy of functions to behave nicely. Lemma 2.14(1,3) is a Ketonen-Solovay counterpart of Bigorajska and Kotlarski [1, Lemma 4]:

Lemma 2.14. *Let h be as above. Let $0 < \alpha, \beta < \epsilon_0$ and $x \in \mathbb{N}$. Then:*

- (1) *h_β is increasing*
- (2) *if $x \in \text{dom } h$ and $h(x) \in \text{dom } h_\beta$ then $x \in \text{dom } h_\beta$*
- (3) *if $\beta \Rightarrow_x \alpha$ and $x \in \text{dom } h_\beta$ then $x \in \text{dom } h_\alpha$ and $h_\alpha(x) \leq h_\beta(x)$*
- (4) *if $x \in \text{dom } h_\beta$, then $h_\beta(x) \geq h(x)$*

Proof. We prove (1),(2) and (3) by mutual induction on β . If $\beta = 1$ then the property holds as $h_1 = h$ is increasing and the only $0 < \alpha < \epsilon_0$ such that $\beta \Rightarrow_x \alpha$ is $\alpha = 1$.

Let $\beta > 1$ and assume that (1), (2) and (3) hold for every $0 < \beta' < \beta$. Let $x, y \in \text{dom } h_\beta$ such that $x < y$, then $h_\beta(y) = h_{\{\beta\}(y)}(h(y))$. As $\{\beta\}(y) < \beta$ and as, by Lemma 2.13(5), $\{\beta\}(y) \Rightarrow_{h(y)} \{\beta\}(x)$, then by (3) for $\{\beta\}(y)$ we have $h(y) \in \text{dom } h_{\{\beta\}(x)}$ and $h_{\{\beta\}(y)}(h(y)) \geq h_{\{\beta\}(x)}(h(y))$. From $x \in \text{dom } h_\beta$ we get $h(x) \in \text{dom } h_{\{\beta\}(x)}$, hence, by (1) for $\{\beta\}(x)$, we get $h_{\{\beta\}(x)}(h(y)) > h_{\{\beta\}(x)}(h(x))$. Thus, $h_\beta(y) > h_\beta(x)$ and h_β is increasing. Thus (1) holds for β .

Let $x \in \text{dom } h$ be such that $h(x) \in \text{dom } h_\beta$. Then we have $h_\beta(h(x)) = h_{\{\beta\}(h(x))}(h(h(x)))$ hence $h(h(x)) \in \text{dom } h_{\{\beta\}(h(x))}$. By (2) for $\{\beta\}(h(x))$ we have $h(x) \in \text{dom } h_{\{\beta\}(h(x))}$. Then, as Lemma 2.13(5) $\{\beta\}(h(x)) \Rightarrow_{h(x)} \{\beta\}(x)$, we also have $h(x) \in \text{dom } h_{\{\beta\}(x)}$ by (3) for $\{\beta\}(h(x))$, hence $x \in \text{dom } h_\beta$. Thus (2) holds for β .

Let $x \in \mathbb{N}$ and $0 < \alpha$ be such that $\beta \Rightarrow_x \alpha$ and $h_\beta(x) \downarrow$. Since $\beta \Rightarrow_x \alpha$, either $\beta = \alpha$ or $\{\beta\}(x) \Rightarrow_x \alpha$. If $\beta = \alpha$ then $h_\alpha(x) \downarrow$ and $h_\alpha(x) = h_\beta(x)$, in which case (3) holds. If $\{\beta\}(x) \Rightarrow_x \alpha$, then as $h_\beta(x) = h_{\{\beta\}(x)}(h(x))$, we have $x \in \text{dom } h_{\{\beta\}(x)}$ by (2) for $\{\beta\}(x)$ and $h_{\{\beta\}(x)}(h(x)) > h_{\{\beta\}(x)}(x)$ by (1) for $\{\beta\}(x)$. From $\{\beta\}(x) \Rightarrow_x \alpha$ and (3) for $\{\beta\}(x)$, we get $x \in \text{dom } h_\alpha$ and $h_{\{\beta\}(x)}(x) \geq h_\alpha(x)$. So $h_\beta(x) = h_{\{\beta\}(x)}(h(x)) > h_{\{\beta\}(x)}(x) \geq h_\alpha(x)$, Thus (3) holds for β .

Last, since $\beta \Rightarrow_x 1$ for every $\beta > 0$ and $x \in \mathbb{N}$, then (4) follows from (3). \square

The following lemma is the Ketonen-Solovay counterpart of Bigorajska and Kotlarski [1, Lemma 7]. Formulated in terms of largeness, it corresponds to Ketonen and Solovay [12, Lemma 4.6].

Lemma 2.15. *Let $\alpha \ll \beta$ and $x \in \mathbb{N}$, if $h_{\beta+\alpha}(x) \downarrow$ then $h_\beta(h_\alpha(x)) \downarrow$ and $h_{\beta+\alpha}(x) = h_\beta(h_\alpha(x))$.*

Proof. By induction on α . If $\alpha = 0$ then, for $x \in \mathbb{N}$ and $\beta \gg \alpha$ such that $h_{\beta+\alpha}(x) \downarrow$, we have $h_\beta(x) \downarrow$, then, as $h_0(x) = x$ we have $h_\beta(x) = h_\beta(h_0(x)) \downarrow$ and $h_\beta(h_0(x)) = h_{\beta+0}(x)$.

Let $\alpha > 0$ and assume the property to hold for every $\alpha' < \alpha$. Let $x \in \mathbb{N}$ and $\beta \gg \alpha$, if $h_{\beta+\alpha}(x) \downarrow$ then $h_{\{\beta+\alpha\}(x)}(h(x)) \downarrow$ and $h_{\{\beta+\alpha\}(x)}(h(x)) = h_{\beta+\alpha}(x)$. By Lemma 2.2, $\{\beta + \alpha\}(x) = \beta + \{\alpha\}(x)$, so $h_{\beta+\{\alpha\}(x)}(h(x)) = h_{\beta+\alpha}(x)$. As $\{\alpha\}(x) < \alpha$ and $\{\alpha\}(x) \ll \beta$, by the induction hypothesis we have $h_\beta(h_{\{\alpha\}(x)}(h(x))) \downarrow$ and $h_\beta(h_{\{\alpha\}(x)}(h(x))) = h_{\beta+\{\alpha\}(x)}(h(x)) = h_{\beta+\alpha}(x)$. And, as $h_{\{\alpha\}(x)}(h(x)) = h_\alpha(x)$, we have $h_{\beta+\alpha}(x) = h_\beta(h_\alpha(x))$. \square

We now prove our core proposition. The intuition goes as follows: Suppose $0 < \lambda < \epsilon_0$ and $x > 0$. The evaluation of $h_\lambda(x)$ unfolds into $h_{\{\lambda\}(x)}(h(x))$, then $h_{\{\{\lambda\}(x)\}(h(x))}(h(h(x)))$, and so on. Thus, the i th iteration of the evaluation of $h_\lambda(x)$, if defined, corresponds to $h_{\lambda_i}(x_i)$, where x_i and λ_i are defined inductively by $x_0 = x$, $\lambda_0 = \lambda$ and $x_{i+1} = h(x_i)$, $\lambda_{i+1} = \{\lambda_i\}(x_i)$. Note that since the function h is increasing, so is the sequence $x_0 < x_1 < \dots$

The definition of $\{\lambda\}(x)$ is done by considering the short Cantor normal form, by reducing first the smallest component of the Cantor normal form. The evaluation either decreases a successor ordinal by one, or replaces a limit ordinal by a smaller one, with coefficient x . Since the sequence $\lambda_0 > \lambda_1 > \dots$ is defined by successive evaluations with an increasing sequence $x_0 < x_1 < \dots$, the higher components will be evaluated with larger values than the smaller components, and therefore this evaluation order ensures that the sequence $\lambda_0 > \lambda_1 > \dots$ decreases as slowly as possible.

Based on this evaluation order, if $\lambda = \beta + \alpha$, with $\beta \gg \alpha$, the sequence $\lambda_0 > \lambda_1 > \dots$ corresponds to evaluating α successively with x_0, x_1, \dots until we reach a stage i such that $\alpha_i = 0$. Then we evaluate β with x_{i+1}, x_{i+2}, \dots

One could consider other evaluation orders, in which one would alternate some evaluation steps of β , and some of α . In terms of the Hardy-like hierarchy, one evaluation step of β corresponds to $h_{\{\beta\}(x) \oplus \alpha}(h(x))$ while one evaluation step of α corresponds to $h_{\beta+\{\alpha\}(x)}(h(x))$. Since the standard evaluation order is the one making the sequence $\lambda_0 > \lambda_1 > \dots$ decrease as slowly as possible, this translates into the following inequality:

$$h_{\beta+\alpha}(x) = h_{\beta+\{\alpha\}(x)}(h(x)) \geq h_{\{\beta\}(x) \oplus \alpha}(h(x))$$

The following proposition formalizes this intuition in the case $\beta \gg \alpha$. It will then be proven in its most general form in Theorem 2.18.

Proposition 2.16. *Let $0 < \alpha, \beta < \epsilon_0$ with $\beta \gg \alpha$ and $x > 0$ an integer. If $h_{\beta+\alpha}(x) \downarrow$, then $h_{\{\beta\}(x) \oplus \alpha}(h(x)) \downarrow$ and $h_{\beta+\alpha}(x) = h_{\beta+\{\alpha\}(x)}(h(x)) \geq h_{\{\beta\}(x) \oplus \alpha}(h(x))$.*

Proof. The proof is by case analysis on the Cantor normal forms, and not by induction. Write $\alpha = \omega^{\alpha_n} \cdot a_n + \dots + \omega^{\alpha_0} \cdot a_0$ and $\beta = \omega^{\beta_k} \cdot b_k + \dots + \omega^{\beta_0} \cdot b_0$ their Cantor normal form. From the assumption $\beta \gg \alpha$, we have $\beta_0 \geq \alpha_n$.

Case 1: $n = 0$ and $\alpha_n = \beta_0$. In that case, $\beta + \{\alpha\}(x) = \{\beta\}(x) \oplus \alpha$, hence $h_{\{\beta\}(x) \oplus \alpha}(h(x)) \downarrow$ and $h_{\beta + \{\alpha\}(x)}(h(x)) = h_{\{\beta\}(x) \oplus \alpha}(h(x))$.

If we are not in this case, we can freely assume that $\alpha_n < \beta_0$ in the Cantor normal form of α and β . Indeed, if $\alpha_n = \beta_0$, we can consider $\alpha' = \omega^{\alpha_{n-1}} a_{n-1} + \dots + \omega^{\alpha_0} a_0$ and $\beta' = \omega^{\beta_k} b_k + \dots + \omega^{\beta_0} (b_0 + a_n)$, we then have $0 < \alpha', \beta' < \epsilon_0$, $\alpha' \ll \beta'$, $\beta + \alpha = \beta' + \alpha'$, $\beta + \{\alpha\}(x) = \beta' + \{\alpha'\}(x)$ and $\{\beta\}(x) \oplus \alpha = \{\beta'\}(x) \oplus \alpha'$, making it sufficient to prove the result for α' and β' .

Case 2: $\alpha_n < \beta_0$ and β_0 is successor. Let β'_0 be such that $\beta_0 = \beta'_0 + 1$. In that case, write $\beta = \delta + \omega^{\beta'_0 + 1}$ and $\alpha = \gamma + \omega^{\alpha_0}$. By our assumption that $\alpha_n < \beta_0$, we have $\omega^{\beta'_0 + 1} > \alpha$ and $\omega^{\beta'_0} \gg \alpha$. By Lemma 2.15, letting $c = h_{\{\alpha\}(x)}(h(x))$, we have

$$h_{\beta + \{\alpha\}(x)}(h(x)) = h_\beta(c) = h_{\delta + \omega^{\beta'_0} \cdot c}(h(c))$$

By Lemma 2.14(4), $c \geq h(x) > x$, so by Lemma 2.13(2), we have $\omega^{\beta'_0} \cdot c \Rightarrow_y \omega^{\beta'_0} \cdot (x + 1)$ for every $y \in \mathbb{N}$, and as $\delta \gg \omega^{\beta'_0} \cdot c$, we have $\delta + \omega^{\beta'_0} \cdot c \Rightarrow_y \delta + \omega^{\beta'_0} \cdot (x + 1)$ for every $y \in \mathbb{N}$ by Lemma 2.13(1). Thus, by Lemma 2.14(3), $h_{\delta + \omega^{\beta'_0} \cdot (x+1)}(h(c)) \downarrow$ and by Lemma 2.15, letting $d = h_{\{\omega^{\beta'_0}\}(h(c))}(h(h(c)))$,

$$h_{\delta + \omega^{\beta'_0} \cdot c}(h(c)) \geq h_{\delta + \omega^{\beta'_0} \cdot (x+1)}(h(c)) = h_{\delta + \omega^{\beta'_0} \cdot x}(h_{\omega^{\beta'_0}}(h(c))) = h_{\delta + \omega^{\beta'_0} \cdot x}(d)$$

By Lemma 2.10, $c \geq \text{psn}(\{\alpha\}(x)) + h(x)$, hence by Lemma 2.5, $c \geq \text{psn}(\alpha)$ and $h(c) > \text{psn}(\{\alpha\}(h(x)))$. By Lemma 2.4, we get $\{\omega^{\beta'_0}\}(h(c)) > \{\alpha\}(h(x))$ from the fact that $\omega^{\beta'_0} > \{\alpha\}(h(x))$ and $h(c) > \text{psn}(\{\alpha\}(h(x)))$, hence, by Lemma 2.12 we have $\{\omega^{\beta'_0}\}(h(c)) \Rightarrow_{h(c)} \{\alpha\}(h(x))$. By Lemma 2.13(1), since $\delta + \omega^{\beta'_0} \cdot x \gg \{\omega^{\beta'_0}\}(h(c))$, we have

$$\delta + \omega^{\beta'_0} \cdot x + \{\omega^{\beta'_0}\}(h(c)) \Rightarrow_{h(c)} \delta + \omega^{\beta'_0} \cdot x + \{\alpha\}(h(x))$$

Thus, by Lemma 2.14(2,3) and Lemma 2.15, $h_{\delta + \omega^{\beta'_0} \cdot x}(h_{\{\alpha\}(h(x))}(h(h(x)))) \downarrow$ and

$$\begin{aligned} h_{\delta + \omega^{\beta'_0} \cdot x}(d) &\geq h_{\delta + \omega^{\beta'_0} \cdot x}(h_{\{\alpha\}(h(x))}(h(h(x)))) \\ &= h_{\delta + \omega^{\beta'_0} \cdot x}(h_\alpha(h(x))) \\ &= h_{\delta + \omega^{\beta'_0} \cdot x + \alpha}(h(x)) \\ &= h_{\{\beta\}(x) \oplus \alpha}(h(x)) \end{aligned}$$

Putting altogether,

$$h_{\beta + \{\alpha\}(x)}(h(x)) = h_{\delta + \omega^{\beta'_0} \cdot c}(h(c)) \geq h_{\delta + \omega^{\beta'_0} \cdot x}(d) \geq h_{\{\beta\}(x) \oplus \alpha}(h(x))$$

Case 3: $\alpha_n < \beta_0$ and β_0 limit. In that case, write $\beta = \delta + \omega^{\beta_0}$ and $\alpha = \gamma + \omega^{\alpha_0}$. Similarly, by Lemma 2.2 and Lemma 2.15, letting $c = h_{\{\alpha\}(x)}(h(x))$, we have

$$h_{\beta+\alpha}(x) = h_{\beta+\{\alpha\}(x)}(h(x)) = h_\beta(c) = h_{\delta+\omega^{\{\beta_0\}(c)}}(h(c))$$

Since $c > x$, we have $\{\beta_0\}(c) \Rightarrow_y \{\beta_0\}(x) + 1$ for every $y \geq 2$ by Lemma 2.13(3,5). Note that $x \geq 1$, hence $h(x) \geq 2$, so by Lemma 2.14(4), $c \geq 2$ and $h(c) \geq 3$. Thus by Lemma 2.13(4),

$$\omega^{\{\beta_0\}(c)} \Rightarrow_{h(c)} \omega^{\{\beta_0\}(x)+1} \Rightarrow_{h(c)} \omega^{\{\beta_0\}(x)} \cdot 3$$

Since $\delta \gg \omega^{\{\beta_0\}(c)}$, by Lemma 2.13(1),

$$\delta + \omega^{\{\beta_0\}(c)} \Rightarrow_{h(c)} \delta + \omega^{\{\beta_0\}(x)} \cdot 3$$

Therefore, by Lemma 2.14(3), $h_{\delta+\omega^{\{\beta_0\}(x)} \cdot 3}(h(c)) \downarrow$ and, letting $d = h_{\omega^{\{\beta_0\}(x)}}(h(c))$,

$$h_{\delta+\omega^{\{\beta_0\}(c)}}(h(c)) \geq h_{\delta+\omega^{\{\beta_0\}(x)} \cdot 3}(h(c)) = h_{\delta+\omega^{\{\beta_0\}(x)} \cdot 2}(d)$$

By Lemma 2.10, $d \geq \text{psn}(\{\omega^{\beta_0}\}(x)) + h(c) \geq \text{psn}(\{\omega^{\beta_0}\}(x)) + 3$ and $d > c \geq \text{psn}(\{\alpha\}(x)) + h(x) \geq \text{psn}(\{\alpha\}(x)) + 2$. Thus, by Lemma 2.5, $d > \text{psn}(\alpha) + 1$ and $d > \max(\text{psn}(\omega^{\{\beta_0\}(x)}), \text{psn}(\alpha) + 1) \geq \text{psn}(\omega^{\{\beta_0\}(x)} \oplus \alpha)$.

From $\omega^{\{\beta_0\}(x)} \cdot 2 > \omega^{\{\beta_0\}(x)} \oplus \alpha$ and Lemma 2.12 we have $\omega^{\{\beta_0\}(x)} \cdot 2 \Rightarrow_d \omega^{\{\beta_0\}(x)} \oplus \alpha$ and by Lemma 2.13(1), $\delta + \omega^{\{\beta_0\}(x)} \cdot 2 \Rightarrow_d \delta + \omega^{\{\beta_0\}(x)} \oplus \alpha$. Finally, by Lemma 2.14(2,3),

$$h_{\delta+\omega^{\{\beta_0\}(x)} \cdot 2}(d) \geq h_{\delta+\omega^{\{\beta_0\}(x)} \oplus \alpha}(h(x)) = h_{\{\beta\}(x) \oplus \alpha}(h(x))$$

This completes our proof of Proposition 2.16. \square

Lemma 2.17. *Let $\alpha, \beta < \epsilon_0$, then there exists $\alpha', \beta' < \epsilon_0$ with $\alpha' \ll \beta'$ and such that $\beta \oplus \alpha = \beta' + \alpha'$ and $\{\beta\}(x) \oplus \alpha = \{\beta'\}(x) \oplus \alpha'$ for every $x \in \mathbb{N}$*

Proof. If $\beta = 0$ then take $\alpha' = \alpha$ and $\beta' = \beta$. If $\beta \neq 0$, then write $\alpha = \omega^{\gamma_n} \cdot a_n + \dots + \omega^{\gamma_0} \cdot a_0$ and $\beta = \omega^{\gamma_n} \cdot b_n + \dots + \omega^{\gamma_0} \cdot b_0$ with $a_0, \dots, a_n, b_0, \dots, b_n \in \mathbb{N}$ and $\gamma_0 < \dots < \gamma_n < \epsilon_0$. Let $i \leq n$ be the smallest index such that $b_i \neq 0$, and let $\beta' = \omega^{\gamma_n} \cdot (b_n + a_n) + \dots + \omega^{\gamma_i} \cdot (b_i + a_i)$ and $\alpha' = \omega^{\gamma_{i-1}} \cdot (b_{i-1} + a_{i-1}) + \dots + \omega^{\gamma_0} \cdot (b_0 + a_0)$.

We have $\beta \oplus \alpha = \beta' + \alpha' = \omega^{\gamma_n} (b_n + a_n) + \dots + \omega^{\gamma_0} (b_0 + a_0)$ and $\{\beta'\}(x) \oplus \alpha' = (\omega^{\gamma_n} (b_n + a_n) + \dots + \omega^{\gamma_i} (b_i + a_i - 1) + \dots + \omega^{\gamma_0} (b_0 + a_0)) \oplus \{\omega^{\gamma_i}\}(x) = \{\beta\}(x) \oplus \alpha$ \square

The following theorem informally states that the evaluation order of the fundamental sequences is the one which makes the Hardy-like hierarchy grow the fastest. In the case $\alpha \ll \beta$, $\beta + \alpha = \beta \oplus \alpha$, hence it generalizes Proposition 2.16.

Theorem 2.18 (Optimal growth). *Let $\alpha, \beta < \epsilon_0$ with $\beta > 0$ and $x > 0$ an integer. If $h_{\beta \oplus \alpha}(x) \downarrow$ then $h_{\{\beta\}(x) \oplus \alpha}(h(x)) \downarrow$ and $h_{\beta \oplus \alpha}(x) = h_{\{\beta \oplus \alpha\}(x)}(h(x)) \geq h_{\{\beta\}(x) \oplus \alpha}(h(x))$.*

Proof. Let $\alpha', \beta' < \epsilon_0$ be obtained from Lemma 2.17 applied on α, β . As $\beta \oplus \alpha = \beta' + \alpha'$ and $\{\beta\}(x) \oplus \alpha = \{\beta'\}(x) \oplus \alpha'$, it is sufficient to show that $h_{\{\beta'\}(x) \oplus \alpha'}(h(x)) \downarrow$ and that $h_{\beta' \oplus \alpha'}(x) = h_{\{\beta' \oplus \alpha'\}(x)}(h(x)) \geq h_{\{\beta'\}(x) \oplus \alpha'}(h(x))$.

If $\alpha' = 0$, then the result holds by definition of $h_{\beta'}$.

If $\alpha' \neq 0$, then the result is that of Proposition 2.16. \square

2.2 Partition theorem

We now translate this abstract study of the Hardy-like hierarchy into the framework of α -largeness, to prove Main Theorem 1.1.

Definition 2.19. Let $A = \{x_0, \dots, x_{n-1}\} \subseteq \mathbb{N}$ a finite set. We define $h^A : A \setminus \{\max A\} \rightarrow A$ by $h^A(x_i) = x_{i+1}$ for every $i < n-1$. In other words, h^A is the successor operation on the ordered set $(A, <_{\mathbb{N}})$.

The following lemma is the heart of the correspondence between Hardy-like hierarchies and of Ketonen and Solovay's α -largeness.

Lemma 2.20. Let $0 < \alpha < \epsilon_0$, then A is at most α -large iff $h_{\alpha}^A(\min A) \uparrow$.

Proof. By induction on α . The result holds for $\alpha = 1$, as $h_{\alpha}^A(\min A) \uparrow$ if and only if $|A| \leq 1$. Let $1 < \alpha < \epsilon_0$ and assume the property holds for every $\beta < \alpha$. Let $A = \{x_0, x_1, \dots, x_n\} \subseteq \mathbb{N}$ be such that $h_{\alpha}^A(x_0) \uparrow$, then, $h_{\{\alpha\}(x_0)}^A(x_1) \uparrow$ and therefore $h_{\{\alpha\}(x_0)}^{A \setminus \{x_0\}}(x_1) \uparrow$. By the inductive hypothesis, $A \setminus \{x_0\}$ is therefore at most $\{\alpha\}(x_0)$ -large, hence A is at most α -large. \square

The following theorem is an iterated counterpart of Theorem 2.18 in the language of α -largeness. It has several consequences, such as the splitting property (Theorem 2.21) and multiple versions of the pigeonhole principle (Theorems 2.22 and 2.23)

Main Theorem 1.1. Let $\beta, \gamma < \epsilon_0$ be two ordinals, and $B, C \subseteq \mathbb{N}$ be at most β -large and γ -large respectively. Then $B \cup C$ is at most $(\beta \oplus \gamma)$ -large.

Proof. By assumption we have $h_{\beta}^B(\min B) \uparrow$ and $h_{\gamma}^C(\min C) \uparrow$. We want to show that $h_{\beta \oplus \gamma}^{B \cup C}(\min\{B, C\}) \uparrow$.

Write $B \cup C = \{x_0 < x_1 < \dots < x_{k-1}\}$ and consider the sequences of ordinals $(\beta_i)_{i < k}$ and $(\gamma_i)_{i < k}$ defined inductively as follows: $\beta_0 = \beta$ and $\gamma_0 = \gamma$ and if β_i and γ_i have been defined, let $\beta_{i+1} = \{\beta_i\}(x_i)$ and $\gamma_{i+1} = \gamma_i$ if $x_i \in B$ and $\beta_{i+1} = \beta_i$ and $\gamma_{i+1} = \{\gamma_i\}(x_i)$ if $x_i \in C$.

Since B is at most β -large, then $\{\beta\}(B \setminus \{\max B\}) > 0$, thus, if $x_i \in B$ then $\beta_i \neq 0$, similarly, if $x_i \in C$ then $\gamma_i \neq 0$. Hence, if $x_{k-1} \in B$ then $\beta_{k-1} \neq 0$ and if $x_{k-1} \in C$ then $\gamma_{k-1} \neq 0$, in every case $\beta_{k-1} \oplus \gamma_{k-1} \neq 0$.

If $h_{\beta_0 \oplus \gamma_0}^{B \cup C}(x_0) \downarrow$, then, by induction on $i < k$, using Theorem 2.18, we claim that for every $i < k$, we have $h_{\beta_i \oplus \gamma_i}^{B \cup C}(x_i) \downarrow$ and $h_{\beta_i \oplus \gamma_i}^{B \cup C}(x_i) \geq h_{\beta_{i+1} \oplus \gamma_{i+1}}^{B \cup C}(x_{i+1})$. Thus, if $h_{\beta_0 \oplus \gamma_0}^{B \cup C}(x_0) \downarrow$ then $h_{\beta_{k-1} \oplus \gamma_{k-1}}^{B \cup C}(x_{k-1}) \downarrow$, which is only possible if $\beta_{k-1} \oplus \gamma_{k-1} = 0$, contradiction. Thus $h_{\beta_0 \oplus \gamma_0}^{B \cup C}(x_0) \uparrow$ and $B \cup C$ is at most $(\beta \oplus \gamma)$ -large. \square

The following splitting property generalizes [12, Lemma 4.6]. Note that the reversal holds if $\beta \gg \alpha$.

Theorem 2.21 (Splitting property). *Let X be $(\alpha \oplus \beta)$ -large for $\alpha, \beta < \epsilon_0$, then there exists $X_0 < X_1 \subseteq X$ such that X_0 is α -large and X_1 is β -large.*

Proof. Let X_0 be the prefix of X that is exactly α -large, i.e., if $X = \{x_0, \dots, x_{k-1}\}$, we let $X_0 = \{x_0, \dots, x_i\}$ for $i < k$ the smallest index such that $\{\alpha\}(x_0, \dots, x_i) = 0$. Such a set X_0 exists, as every coefficient in the Cantor normal form of $\alpha \oplus \beta$ is bigger than the corresponding one in the Cantor normal form of α .

Let $X_1 = X \setminus X_0$, $\star = \max X + 1$ and let $X' = X \cup \{\star\}$. By the contrapositive of Main Theorem 1.1, since $X' = X_0 \cup (X_1 \cup \{\star\})$ is not at most $(\alpha \oplus \beta)$ -large, either X_0 is not at most α -large or $X_1 \cup \{\star\}$ is not at most β -large. Thus, $X_1 \cup \{\star\}$ is not at most β -large, and X_1 is therefore β -large. \square

We now deduce two versions of the pigeonhole principle.

Theorem 2.22 (First pigeonhole principle). *Let $X = X_0 \cup X_1$ be $(\alpha \oplus \beta)$ -large for some $\alpha, \beta < \epsilon_0$, then either $X_0 \setminus \{\max X_0\}$ is α -large or $X_1 \setminus \{\max X_1\}$ is β -large or X_0 and X_1 are exactly α -large and exactly β -large, respectively.*

Proof. Suppose X_0 is at most α -large, otherwise $X_0 \setminus \{\max X_0\}$ is α -large and we are done. Similarly, suppose X_1 is at most β -large, otherwise $X_1 \setminus \{\max X_1\}$ is β -large and we are done. Let $\star = \max X + 1$. The set $X_0 \cup (X_1 \cup \{\star\})$ is not at most $(\alpha \oplus \beta)$ -large, hence, by the contrapositive of Main Theorem 1.1, $X_1 \cup \{\star\}$ is not at most β -large. Since X_1 is at most β -large, it is exactly β -large. By applying the contrapositive of Main Theorem 1.1 on the partition $(X_0 \cup \{\star\}) \cup X_1$, we get that $X_0 \cup \{\star\}$ is not at most α -large, hence since X_0 is at most α -large, X_0 is exactly α -large and we are also done since both X_0 and X_1 are exactly β -large. \square

Theorem 2.23 (Second pigeonhole principle). *Let $X = X_0 \cup X_1 \cup \{\star\}$ be $(\alpha \oplus \beta)$ -large for some $\alpha, \beta < \epsilon_0$ and some $\star > X_0 \cup X_1$. Then either X_0 is α -large or X_1 is β -large.*

Proof. Let $\star' = \max X + 1$. The set $X \cup \{\star'\} = (X_0 \cup \{\star\}) \cup (X_1 \cup \{\star'\})$ is not at most $(\alpha \oplus \beta)$ -large, hence, by the contrapositive of Main Theorem 1.1, either $X_0 \cup \{\star\}$ is not at most α -large, in which case X_0 is α -large, or $X_1 \cup \{\star'\}$ is not at most β -large, in which case X_1 is β -large. \square

Since α -largeness is closed under supersets, if $X_0 \cup X_1$ is $(\alpha \oplus \beta)$ -large, then so is $X_0 \cup X_1 \cup \{\star\}$ for any $\star > \max X_1$, in which case either X_0 is α -large or X_1 is β -large. This formulation of the pigeonhole principle, although more natural, is less optimal than Theorem 2.23.

3 Largeness below ω^ω

We now turn to the study of α -largeness in the restricted setting of $\alpha < \omega^\omega$. We shall give a particular focus to ordinals of the form $\omega^n \cdot k$ for $n, k \geq 1$. Thanks to Ketonen and Solovay [12, Lemma 4.6], there exists a simple inductive definition of α -largeness for ordinals of this form:

Proposition 3.1 (Ketonen and Solovay [12]). *A set $F \subseteq_{\text{fin}} \mathbb{N}$ is*

- (1) ω^0 -large iff $F \neq \emptyset$
- (2) $\omega^{(n+1)}$ -large iff $F \setminus \min F$ is $(\omega^n \cdot \min F)$ -large
- (3) $\omega^n \cdot k$ -large iff there are k ω^n -large subsets of F

$$F_0 < F_1 < \dots < F_{k-1}$$

Proof. (1) and (2) are simply unfolding of the definition. (3) is by induction on k : The case $k = 1$ is trivial. Assuming (3) holds for k , by Ketonen and Solovay [12, Lemma 4.6], a set X is $\omega^n \cdot (k+1)$ -large iff there exist two sets $X_0 < X_1 \subseteq X$ such that X_0 is ω^n -large and X_1 is $\omega^n \cdot k$ -large. Apply the induction hypothesis on X_1 . \square

3.1 Construction and deconstruction

Thanks to Proposition 3.1, any ω^{n+1} -large set X can be seen as an increasing sequence of $\min X$ many ω^n -large blocks. In this section, we prove two important propositions: a *construction* property, which quantifies how many ω^n -large blocks a sufficient to produce an ω^{n+m} -large set, and a *deconstruction* property, which counts the number of ω^n -large blocks which can be extracted from an ω^{n+m} -large set.

The quantification is expressed in terms of α -largeness. There are multiple ways to give a meaning to the sentence “the sequence of blocks $X_0 < \dots < X_{d-1}$ is α -large”. One could ask that $\{\max X_i : i < d\}$ is α -large, or that any sequence in $X_0 \times \dots \times X_{d-1}$ is α -large. Thanks to the regularity of α -largeness ([12, Lemma 2]), the two definitions are equivalent.

Proposition 3.2 (Construction). *For every $n, m, d \in \mathbb{N}$ and for every sequence of ω^n -large sets $X_0 < \dots < X_{d-1}$ such that $\{\max X_i : i < d\}$ is ω^m -large, then $X_0 \cup \dots \cup X_{d-1}$ is ω^{n+m} -large.*

Proof. By induction on m . The case $m = 0$ is immediate, since any ω^0 -large set is non-empty, so $d \geq 1$ and X_0 is ω^n -large. Suppose it holds for m . If $Y = \{\max X_i : i < d\}$ is ω^{m+1} -large, then $Y \setminus \{\min Y\}$ is $\omega^m \cdot \min Y$ -large. By Proposition 3.1, there are $\min Y$ many ω^m -large subsets $Y_0 < \dots < Y_{\min Y-1}$ of $Y \setminus \{\min Y\}$. For every $i < \min Y$, let $Z_i = \bigcup_{j \in Y_i} X_j$. By induction hypothesis, each Z_i is ω^{n+m} -large, so $\{\min Y\} \cup Z_0 \cup \dots \cup Z_{\min Y-1}$ is ω^{n+m+1} -large. By upward-closure of α -largeness, $X_0 \cup \dots \cup X_{d-1}$ is ω^{n+m+1} -large. \square

The following deconstruction proposition is an adaptation of Kołodziejczyk and Yokoyama [14, Lemma 2.1]. Note that the hypothesis requires an extra ω^n -large set, while the conclusion also contains some extra ω^n -large set. This might seem sub-optimal but the ω^n -large set of the hypothesis is on the left part of X , while the one from the conclusion is on the right, which is weaker.

Proposition 3.3 (Deconstruction). *For every $n, m \in \mathbb{N}$ and for every $\omega^{n+m} + \omega^n$ -large set X , there are some $d \in \mathbb{N}$ and some ω^n -large subsets $X_0 < \dots < X_d$ of X such that $\{\max X_i : i < d\}$ is ω^m -large.*

Proof. By induction on m . The case $m = 0$ is immediate by Ketonen and Solovay [12, Lemma 4.6], as X is $\omega^n \cdot 2$ -large, so there are two ω^n -large sets $X_0 < X_1 \subseteq X$. Note that $\{\max X_0\}$ is non-empty, hence ω^0 -large.

Suppose it holds for m . Let X be $\omega^{n+m+1} + \omega^n$ -large. By Ketonen and Solovay [12, Lemma 4.6] (or by Theorem 2.21), there are some $A < B \subseteq X$ such that A is ω^n -large and B is ω^{n+m+1} -large. By Proposition 3.1, there are $\min B$ many ω^{n+m} -large sets $Z_0 < \dots < Z_{\min B-1} \subseteq B \setminus \{\min B\}$.

Let $d_{-1} = 0$ and $X_{d_{-1}}^{-1} = Z_0$. Note that $X_{d_{-1}}^{-1}$ is ω^{n+m} -large, hence ω^n -large. We are going to build inductively for every $i < \min B - 1$ a family of ω^n -large sets $X_0^i < \dots < X_{d_i}^i$ such that

$$(1) \quad X_0^i \cup \dots \cup X_{d_i}^i \subseteq X_{d_{i-1}}^{-1} \cup Z_i;$$

$$(2) \quad \{\max X_j^i : j < d_i\} \text{ is } \omega^m\text{-large.}$$

Assuming $X_{d_{i-1}}^{-1} < Z_i$ is defined, the set $X_{d_{i-1}}^{-1} \cup Z_i$ is $\omega^{n+m} + \omega^n$ -large, so by induction hypothesis, there is a sequence $X_0^i, \dots, X_{d_i}^i$ of ω^n -large sets satisfying (1) and (2). It follows that $\{\max X_j^i : i < \min B - 1, j < d_i\}$ is $\omega^m \cdot (\min B - 1)$ -large. Since $\max A < \min B$, then $\{\max A\} \cup \{\max X_j^i : i < \min B - 1, j < d_i\}$ is ω^{m+1} -large, so altogether, these ω^n -large sets form the desired sequence. \square

Corollary 3.4 (Generalized deconstruction). *For every $n, k, \ell \geq 1$ and $m \geq 0$, for every $\omega^{n+m} \cdot (k\ell + 1)$ -large set X such that $k(\ell + 1) \leq \min X$, there are some $d \in \mathbb{N}$ and some $\omega^n \cdot k$ -large subsets $X_0 < \dots < X_{d-1}$ of X such that $\{\max X_i : i < d\}$ is $\omega^m \cdot \ell$ -large.*

Proof. Suppose $m = 0$. Then X is $\omega^n \cdot (k\ell + 1)$ -large, and a fortiori $\omega^n \cdot k\ell$ -large hence there is a sequence $X_0 < \dots < X_{k\ell-1}$ of ω^n -large subsets of X . For every $i < \ell$, let $Z_i = X_{ki} \cup \dots \cup X_{k(i+1)-1}$. Each Z_i is $\omega^n \cdot k$ -large, and $\{\max Z_i : i < \ell\}$ is ℓ -large.

Suppose $m > 0$. In particular, X is $\omega^{n+m} \cdot k\ell + \omega^{n+1}$ -large. Since $k(\ell + 1) \leq \min X$, then X is $\omega^{n+m} \cdot k\ell + \omega^n \cdot k(\ell + 1)$ -large. By Theorem 2.21, there is a sequence $X_{-1} < X_0 < \dots < X_{k\ell-1}$ of subsets of X such that X_{-1} is $\omega^n \cdot k$ -large and $X_0, \dots, X_{k\ell-1}$ are $\omega^{n+m} + \omega^n$ -large. By Proposition 3.3, for each i such that $0 \leq i < k\ell$, there is a sequence $X_i^0 < \dots < X_i^{d_i-1}$ of ω^n -large subsets of X_i such that $\{\max X_j^i : j < d_i\}$ is ω^m -large. Let W_0, \dots, W_{p-1} be the sequence $(X_i^j)_{i < k\ell, j < d_i}$. In particular, the set $Z = \{\max W_i : i < p\}$ is $\omega^m \cdot k\ell$ -large. Let

$g : Z \rightarrow k$ be such that $g(x)$ is the remainder of the euclidean division of i by k , where $x = \max W_i$. By Theorem 2.23, there is an $\omega^m \cdot \ell$ -large g -homogeneous subset $H = \{x_0 < \dots < x_{d-1}\} \subseteq Z$. For each $s < d$, let i_s be such that $\max W_{i_s} = x_s$. Let $H_0 = X_{-1} \cup W_{i_0}$ and $H_{s+1} = \bigcup_{i_s < j \leq i_{s+1}} W_j$. Note that for each $s < d$, $\max H_s = x_s$, so $\{\max H_s : s < d\}$ is $\omega^m \cdot \ell$ -large. Since X_{-1} is $\omega^n \cdot k$ -large, so is H_0 . By g -homogeneity of H , $i_{s+1} \geq i_s + k$, so H_{s+1} is $\omega^n \cdot k$ -large. \square

3.2 Sparsity

The notion of α -largeness induces a dual notion of α -sparsity as follows:

Definition 3.5. A set $X \subseteq \mathbb{N}$ is α -sparse if for every $x, y \in X$ such that $x < y$, $(x, y]_{\mathbb{N}}$ is α -large.

Theorem 3.6 (Ketonen and Solovay [12]). For every primitive recursive function $g : \mathbb{N} \rightarrow \mathbb{N}$, there is some $n \in \mathbb{N}$ such that for every ω^n -large set F , $\max F > g(\min F)$.

In particular, $y > 2x$ whenever $(x, y]$ is ω -large and $y > x2^x$ whenever $(x, y]$ is ω^2 -large. We shall be particularly interested in $\omega^2 \cdot 3$ -sparse sets as it provides sufficient sparsity to bound Ramsey numbers.

Let $R_k(d)$ be the least number R such that for every coloring $f : [R]^2 \rightarrow k$, there is an f -homogeneous set of size d . The standard upper bound for the finite Ramsey's theorem for pairs and two colors, obtained by Erdős and Szekeres in [8], is that $R_2(d) \leq \binom{2d-1}{d-1} \leq 4^d$. The upper bound generalizes to $R_k(d) \leq k^{kd}$ (see Graham, Rothschild and Spencer [9, Section 1.1]).

Lemma 3.7. If X is $\omega^2 \cdot 3$ -sparse and $\min X \geq 7$, then it is $(x \mapsto x^{R_x(2x+2)})$ -sparse.

Proof. Any ω^2 -sparse set is $(x \mapsto x2^x)$ -sparse, so any $\omega^2 \cdot 3$ -sparse set is $(x \mapsto x2^{x+x2^x+x2^{x+x2^x}})$ -sparse, and in particular $(x \mapsto 2^{x2^{x2^x}})$ -sparse. Whenever $x \geq 7$, then $x(2x+2) \leq 2^x$, so $2^{x2^{x2^x}} \geq x^{2^{x^2(2x+2)}} \geq x^{x^{x(2x+2)}}$. Thus X is $(x \mapsto x^{x^{x(2x+2)}})$ -sparse.

Since $R_x(2x+2) \leq x^{x(2x+2)}$ (see Graham, Rothschild and Spencer [9, Section 1.1]), $x^{R_x(2x+2)} \leq x^{x^{x^{x(2x+2)}}}$, so X is $(x \mapsto x^{R_x(2x+2)})$ -sparse. \square

Thanks to the deconstruction proposition Corollary 3.4, one can relate largeness to sparsity. The following lemma specializes this relation in the case of $\omega^2 \cdot 3$ -sparsity.

Lemma 3.8. For every $\ell \geq 1$ and $m \geq 0$, for every ω^{m+3} -large set X such that $3\ell + 2 \leq \min X$, there is some $\omega^2 \cdot 3$ -sparse $\omega^m \cdot \ell + 1$ -large subset of X .

Proof. Since $3\ell + 2 \leq \min X$, then $X \setminus \{\min X\}$ is $\omega^{m+2} \cdot (3\ell + 2)$ -large, hence $\omega^{m+2} \cdot (3\ell + 1) + \omega^{m+2}$ -large. Then there are two subsets $A < B \subseteq X$ such that A is ω^{m+2} -large and B is $\omega^{m+2} \cdot (3\ell + 1)$ -large. Since $3\ell + 2 \leq \min A$,

then $3(\ell + 1) \leq \max B$, so by Corollary 3.4 applied to B , there is a sequence $X_0 < \dots < X_{d-1}$ of $\omega^2 \cdot 3$ -large subsets of B such that $Z = \{\max X_i : i < d\}$ is $\omega^m \cdot \ell$ -large. The set $\{\min X\} \cup Z$ is $\omega^m \cdot \ell + 1$ -large, and $\omega^2 \cdot 3$ -sparse. \square

3.3 Largeness for combinatorial theorems

We are particularly interested in the closure of notions of largeness under combinatorial statements coming from Ramsey theory. Given a coloring $f : [\mathbb{N}]^n \rightarrow k$, a set $H \subseteq \mathbb{N}$ is *f-homogeneous* if f is constant on $[H]^n$.

Statement (Ramsey's theorem). *Given $n, k \in \mathbb{N}$, RT_k^n is the statement “For every coloring $f : [\mathbb{N}]^n \rightarrow k$, there is an infinite f -homogeneous set”.*

Ramsey's theorem for pairs is a central subject of study in reverse mathematics, a foundational program whose goal is to find optimal axioms to prove ordinary theorems [6, 25]. Its first-order consequences are closely related to the computation of bounds for its largeness counterpart:

Definition 3.9. *A set $X \subseteq_{\text{fin}} \mathbb{N}$ is RT_k^n - α -large if for every coloring $f : [X]^n \rightarrow k$, there is an α -large f -homogeneous subset $H \subseteq X$.*

Whenever a parameter is omitted, it is taken to be $\min X$. For instance, a set X is RT^n - α -large if it is $\text{RT}_{\min X}^n$ - α -large, in other words, if for every coloring $f : [X]^n \rightarrow \min X$, there is an α -large f -homogeneous subset $H \subseteq X$. Bounds largeness for Ramsey's theorem for pairs were extensively studied. Ketomänen and Solovay [12] proved that every ω^{k+4} -large is RT_k^2 - ω -large set when $k \geq 2$. Patey and Yokoyama [22] proved that for every $k \in \mathbb{N}$, there is some $n \in \mathbb{N}$ such that every ω^n -large set is RT_2^2 - ω^n -large. Kołodziejczyk and Yokoyama [14] computed some explicit bounds, a proved that every $\omega^{144(n+1)}$ -large set is RT_2^2 - ω^n -large. Our goal is to give a tight bound to RT_k^2 - ω^n -largeness, up to an additive constant.

In order to better understand its computational and proof-theoretic strength, Bovykin and Weiermann [4] decomposed Ramsey's theorem for pairs and two colors (RT_2^2) into two statements: the Erdős-Moser theorem (**EM**) and the Ascending Descending Sequence (**ADS**). The Erdős-Moser theorem [7] is a statement from graph theory about tournaments, that is, complete oriented graphs, which says that any infinite tournaments admits an infinite transitive sub-tournament. The Ascending Descending Sequence principle [11] is a theorem from order theory, stating that every infinite linear order admits an infinite ascending or descending sequence. Both statements can be formulated as particular cases of Ramsey's theorem for pairs, thanks to the notion of transitivity.

Definition 3.10. *Given a coloring $f : [\mathbb{N}]^2 \rightarrow k$, a set $H \subseteq \mathbb{N}$ is f -transitive if for every $x < y < z \in H$ such that $f(x, y) = f(y, z)$, then $f(x, z) = (x, z)$. Whenever $H = \mathbb{N}$, we simply say that the coloring f is transitive.*

Tournaments are in one-to-one correspondence with 2-colorings of pairs, and transitive tournaments with transitive colorings. Similarly, linear orders can be

coded as transitive 2-colorings of pairs, in which case ascending or descending sequences correspond to homogeneous sets. Therefore, given any instance of Ramsey's theorem for pairs and two colors, one can apply **EM** to obtain an infinite transitive subset, and then **ADS** to obtain a homogeneous subset. Both statements were extensively studied in computability theory and reverse mathematics [4, 20, 11, 19].

When trying to extend the decomposition of RT_2^2 into **EM** and **ADS** to more colors, there are two natural generalizations of the notion of transitivity: a weak one, given in Definition 3.10, and a stronger one, formally introduced by Towsner and Yokoyama [26], but already used in Patey [21, Theorem 8.2.1].

Definition 3.11. *Given a coloring $f : [\mathbb{N}]^2 \rightarrow k$, a set $H \subseteq \mathbb{N}$ is f -fallow if for every $x < y < z \in H$, $f(x, z) \in \{f(x, y), f(y, z)\}$.*

Clearly, any f -fallow set is f -transitive, but the converse is not true. Both **EM** and **ADS** therefore admit two generalized statements, based on transitivity and fallowness. We shall only consider the stronger version of each statements, that is, the fallow version of the Erdős-Moser theorem and transitive Ramsey's theorem for pairs.

Statement (Fallow Erdős-Moser theorem). *Given $k \in \mathbb{N}$, fEM_k is the statement “For every coloring $f : [\mathbb{N}]^2 \rightarrow k$, there is an infinite f -fallow set”.*

Statement (Transitive Ramsey's theorem for pairs). *Given $k \in \mathbb{N}$, trRT_k^2 is the statement “For every transitive coloring $f : [\mathbb{N}]^2 \rightarrow k$, there is an infinite f -homogeneous set”.*

The statements fEM_k and trRT_k^2 admit largeness counterparts, as Ramsey's theorem. In the remainder of this article, we shall compute bounds for fEM - ω^n -largeness and trRT_k^2 - ω^n -largeness, and deduce bounds for RT_k^2 - ω^n -largeness.

3.4 Pigeonhole principle

We already studied α -large versions of the pigeonhole principle for $\alpha < \epsilon_0$ in Section 2. The following proposition is an immediate consequence of Theorem 2.23, restricted to the setting of ω^n -largeness. It is however useful to state it under this form as it will be used all over the remainder of the article. This strengthens Kołodziejczyk and Yokoyama [14, Lemma 2.2] by improving the bounds and removing the sparsity assumptions.

Proposition 3.12. *Fix $n \geq 0$ and $a, k \geq 1$.*

- (1) *If X is $\omega^n \cdot ak$ -large, then it is RT_k^1 - $\omega^n \cdot a$ -large.*
- (2) *If X is ω^{n+1} -large, then it is RT^1 - ω^n -large.*

Proof. (1) By induction on k . The case $k = 1$ is trivial. Suppose it holds for k . Let X be $\omega^n \cdot a(k+1)$ -large and $(x \mapsto 2x+1)$ -sparse, and let $f : X \rightarrow k+1$. Let $Y_0 = \{x \in X : f(x) < k\}$ and $Y_1 = \{x \in X : f(x) = k\}$. By Theorem 2.23,

either Y_0 is $\omega^n \cdot ak$ -large, and which case by induction hypothesis, there is an f -homogeneous set which is $\omega^n \cdot a$ -large, or Y_1 is $\omega^n \cdot a$ -large, and is f -homogeneous for color k by definition.

(2) Let $f : X \rightarrow \min X$. Since X is ω^{n+1} -large, $X \setminus \{\min X\}$ is $\omega^n \cdot \min X$ -large. By (1), $X \setminus \{\min X\}$ is $\text{RT}_{\min X - \omega^n}$ -large, so there is an ω^n -large f -homogeneous subset. \square

3.5 Grouping principle

The inductive proofs of Kołodziejczyk and Yokoyama for largeness for EM heavily depended on the notion of grouping. We shall only us the following lemma, which, in the terminology of grouping, means that every $\omega^n \cdot R_2(2d)$ -large $(x \mapsto x2^x)$ -sparse set admits an $(\omega^n + 1, d)$ -grouping. This is an improvement of their bound, which used an ω^{n+3} -large set [14, Lemma 2.6].

Lemma 3.13. *Let $k > 0$ and let $X \subseteq_{fin} \mathbb{N}$ be $\omega^n \cdot R_k(2d)$ -large and $(x \mapsto xk^x)$ -sparse, for some $d \in \mathbb{N}$ such that $k^{R_k(2d)} \leq \min X$ and let $f : [X]^2 \rightarrow k$ be a coloring. Then, there exists a sequence $Y_0 < \dots < Y_{d-1}$ of $\omega^n + 1$ -large subsets of X and some color $c < k$ such that $f(x, y) = c$ for every $x \in Y_s$ and $y \in Y_t$ for $s < t < d$.*

Proof. If $n = 0$, the results holds by definition of R and ω^0 -largeness. Indeed, let $f : [X]^2 \rightarrow k$ be a coloring. Since $|X| \geq R_k(2d)$, by finite Ramsey's theorem for pairs, there is an f -homogeneous set $\{y_0 < y_1 < \dots < y_{2d-1}\} \subseteq X$. Since any singleton element is ω^0 -large, the set $Y_i = \{y_{2i}, y_{2i+1}\}$ is $\omega^0 + 1$ -large. Assume from now on that $n > 0$.

Let $X_0 < \dots < X_{R_k(2d)-1}$ be a decomposition of X into ω^n -large sets. Consider the sequence $(f_i)_{i < R_k(2d)}$ of colorings and the sequence $Z_0 < \dots < Z_{R_k(2d)-1}$ of sets defined inductively as follows:

Assume that $Z_{i+1}, \dots, Z_{R_k(2d)-1}$ have been defined for some $i < R_k(2d)$, and consider the coloring $f_i : X_i \rightarrow k^{R_k(2d)-1-i+\sum_{j < i} |X_j|}$ defined by

$$f_i(x) = (f(y, x))_{y \in \bigcup_{j < i} X_j} \sqcup (f(x, y))_{y \in \{\min Z_j : i < j < R_k(2d)\}}$$

As $\min X_0 \geq R_k(2d)$ and as the X_i are disjoint, we get that for $i > 0$, $R_k(2d) + \sum_{j < i} |X_j| \leq \max X_{i-1}$. By $(x \mapsto xk^x)$ -sparsity of X , $k^{\max X_{i-1}} \cdot \max X_{i-1} \leq \min X_i$, so $k^{R_k(2d) + \sum_{j < i} |X_j|} \cdot \max X_{i-1} \leq \min X_i$. By Proposition 3.12(1), there exists some $\omega^{n-1} \cdot (\max X_{i-1})$ -large and f_i -homogeneous subset $Z_i \subseteq X_i$. For $i = 0$, since $k^{R_2(2d)} \leq \min X_0$, by Proposition 3.12(1), there exists some ω^{n-1} -large and f_0 -homogeneous subset $Z_0 \subseteq X_0$.

The $(Z_i)_{i < R_k(2d)}$ are defined such that, for every $i < j < R_k(2d)$ and $x \in Z_i$, $y \in Z_j$, we have $f(x, y) = f(\min Z_i, \min Z_j)$. By definition of $R_k(2d)$, there exists some subset $\{i_0 < \dots < i_{2d-1}\} \subseteq \{0, \dots, R_k(2d) - 1\}$ and some color $c < k$ such that, for every $j < k < 2d$, $x \in Z_{i_j}$ and $y \in Z_{i_k}$, $f(x, y) = c$.

Finally, let $Y_j = \{z_{2j}^0, z_{2j}^1\} \cup Z_{2j+1}$ for every $j < d$, where z_{2j}^0 and z_{2j}^1 are the two last elements of Z_{2j} . Every Y_j is $\omega^n + 1$ -large. \square

3.6 Erdős-Moser theorem

The goal of this section is to prove that every ω^{n+3} -large set X with $\min X \geq 5$ is $\text{EM-}\omega^n$ -large. The previous known bound of Kołodziejczyk and Yokoyama [14] was ω^{36n+4} -largeness.

Proposition 3.14. *Let $n \geq 0$, let $X \subseteq_{\text{fin}} \mathbb{N}$ be and $(x \mapsto x^{R_x(2x)})$ -sparse,*

- (1) *If X is $\omega^n + 1$ -large with $2 \leq \min X$, then it is $\text{fEM-}\omega^n$ -large.*
- (2) *If X is $\omega^n \cdot R_k(2d)$ -large with $k^{R_k(2d)} \leq \min X$, then it is $\text{fEM}_k\text{-}\omega^n \cdot d$ -large.*
- (3) *If X is $\omega^n \cdot 4$ -large, then it is $\text{fEM-}\omega^n + 1$ -large.*

Proof. We prove (1) and (2) by mutual induction on n . (1) holds trivially for $n = 0$, as any ω^0 -large set is non-empty, and any singleton is $\text{EM-}\omega^0$ -large.

Suppose (1) holds for n . Let us show that (2) holds for n . Let $k, d \in \mathbb{N}$, let $X \subseteq_{\text{fin}} \mathbb{N}$ be $(x \mapsto x^{R_x(2x)})$ -sparse, $(\omega^n \cdot R_k(2d))$ -large and such that $k^{R_k(2d)} \leq \min X$ and fix some coloring $f : [X]^2 \rightarrow k$. As $k^{R_k(2d)} \leq \min X$, we have $k \leq \min X$, hence X is $(x \mapsto xk^x)$ -sparse, and by Lemma 3.13, there exists some sequence $Z_0 < \dots < Z_{d-1}$ of $\omega^n + 1$ -large subsets of X and some color $c < k$ such that $f(x, y) = c$ for every $x \in Z_s$ and $y \in Z_t$ for $s < t < d$. By (1) for n , for each $i < d$, there is an f -fallow ω^n -large subset $W_i \subseteq Z_i$. Then, the set $W = \bigcup_{i < d} W_i$ is $\omega^n \cdot d$ -large and f -fallow.

Indeed, pick by contradiction some $x < y < z \in W$ such that $f(x, z) \notin \{f(x, y), f(y, z)\}$. We cannot have $Z_i \cap \{x, y, z\} = \{x\}$ or $Z_i \cap \{x, y, z\} = \{z\}$ for some $i < d$ as it would imply $f(x, y) = f(x, z) = c$ or $f(x, z) = f(y, z) = c$ by our application of Lemma 3.13. Similarly we cannot have $Z_i \cap \{x, y, z\} = \{y\}$ for some $i < d$ as it implies the existence of some $j < d$ such that $Z_j \cap \{x, y, z\} = \{z\}$, since $x < y < z$. And finally, we cannot have $Z_i \cap \{x, y, z\} = \{x, y, z\}$ for some $i < d$ by our application of the inductive hypothesis.

Suppose (2) holds for n . Let us show (1) holds for $n + 1$. Let $X \subseteq_{\text{fin}} \mathbb{N}$ be $(x \mapsto x^{R_x(2x)})$ -sparse and $(\omega^{n+1} + 1)$ -large and let $x_0 = \min X$. Fix some coloring $f : [X]^2 \rightarrow x_0$, then, $X \setminus \{x_0\}$ is $\omega^n \cdot 2R_{x_0}(2x_0)$ -large as $2R_{x_0}(2x_0) \leq x_0^{R_{x_0}(2x_0)}$ when $x_0 \geq 2$ and $x_0^{R_{x_0}(2x_0)} \leq \min X \setminus \{x_0\}$. By Proposition 3.12(1), there exists some $\omega^n \cdot R_{x_0}(2x_0)$ -large subset $Y \subseteq X \setminus \{x_0\}$ such that $f(x_0, y) = f(x_0, z)$ for every $y, z \in Y$. By (2) for n , there is some f -fallow $\omega^n \cdot x_0$ -large subset $Z \subseteq Y$. Then, the set $W = \{x_0\} \cup Z$ is f -fallow and ω^{n+1} -large.

Indeed, pick by contradiction some $x < y < z \in W$ such that $f(x, z) \notin \{f(x, y), f(y, z)\}$. We cannot have $x = x_0$ as $f(x_0, y) = f(x_0, z)$ for every $y, z \in Y$. And we cannot have $x_0 \in Z$, otherwise $\{x, y, z\} \subseteq Z$, contradicting the fact that Z is f -fallow.

(3) For $n = 0$, if X is $\omega^0 \cdot 4$ -large, then $\text{card } X \geq 4$. In particular, any two-element subset of X is $\text{fEM-}\omega^0 + 1$ -large. Suppose $n > 0$. Fix a coloring $f : [\mathbb{N}]^2 \rightarrow \min X$. Let $x_0 = \min X$ and $g : X \rightarrow x_0$ be defined by $g(x_0) = 0$

and $g(y) = f(x_0, y)$ for $y \neq x_0$. By Proposition 3.12(1), there is an $\omega^n \cdot 2$ -large g -homogeneous subset $Y \subseteq X$. Let $Y_0 < Y_1$ be its decomposition into ω^n -large sets. Let $x_1 = \max Y_0$. Since $n > 0$, $\text{card } Y_0 \geq 2$, so $x_1 > x_0$ and $\{x_1\} \cup Y_1$ is $\omega^n + 1$ -large. By (1), there is a ω^n -large f -fallow subset $Z \subseteq \{x_1\} \cup Y_1$. Since Z is g -homogeneous, then $\{x_0\} \cup Z$ is f -fallow, and $\omega^n + 1$ -large. \square

Corollary 3.15. *Let $n \geq 0$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{n+3} -large and $\min X \geq 7$, then it is $\text{fEM-}\omega^n$ -large.*

Proof. Since $\min X \geq 5$, then by Lemma 3.8, there is an $\omega^2 \cdot 3$ -sparse $\omega^n + 1$ -large subset $Y \subseteq X$. By Lemma 3.7, since $\min X \geq 7$, Y is $(x \mapsto x^{R_x(2x+2)})$ -sparse, so by Proposition 3.14, Y is $\text{fEM-}\omega^n$ -large. \square

3.7 Transitive colorings

The strategy to compute bounds for transitive Ramsey's theorem for pairs and k -colors is different from the one for fallow Erdős-Moser theorem. Patey and Yokoyama [22, Lemma 4.4] proved that every ω^{2n+6} -large set is ADS- ω^n -large, using a slightly modified notion of largeness. The proof consisted in defining a coloring $f : [\mathbb{N}]^2 \rightarrow 2n+2$ and applying Ketonen and Solovay's bound [12]. Kołodziejczyk and Yokoyama [14, Theorem 2.11] gave a more direct but less optimal proof by getting rid of the intermediate notion of largeness, and showed that any ω^{4n+4} -large set X with $\min X \geq 3$ is $\text{trRT}_2^2\text{-}\omega^n$ -large.

In this section, we give a direct and optimal proof, by generalizing the bound of Ketonen and Solovay to ω - g -largeness. Given an increasing function $g : \mathbb{N} \rightarrow \mathbb{N}$, a set X is ω - g -large if $\text{card } X > g(\min X)$. Whenever g is the identity function, this yields the standard notion of ω -largeness. The following proposition is a generalization of Ketonen and Solovay [12, Lemma 6.3] and of Kołodziejczyk and Yokoyama [14, Theorem 1.5]:

Proposition 3.16. *Fix a primitive recursive function $g : \mathbb{N} \rightarrow \mathbb{N}$. If X is $(\omega^k + 1)$ -large and g -sparse for $k = 1$ and $(x \mapsto k^{R_k(2g(x))})$ -sparse for $k \geq 2$, then it is $\text{RT}_k^2\text{-}\omega$ - g -large.*

Proof. By induction on k . Suppose $k = 1$. Let $f : X \rightarrow 1$ be the trivial constant coloring. Then X is f -homogeneous. Since X is $\omega^k + 1$ -large, then, letting $X = \{x_0 < x_1\} \sqcup Y$, then $\text{card } Y \geq x_1$ and by g -sparsity of X , $x_1 > g(x_0)$, so $\text{card } X > g(\min X)$.

Assume the property to be true for some $k > 0$. Let X be $(\omega^{k+1} + 1)$ -large and $(x \mapsto (k+1)^{R_{k+1}(2g(x))})$ -sparse and consider a coloring $f : [X]^2 \rightarrow k+1$. Let $x_0 = \min X$, by sparsity, $\min(X \setminus \{x_0\}) \geq (k+1)^{R_{k+1}(2g(x_0))}$ and $X \setminus \{x_0\}$ is $(\omega^k \cdot (k+1) \times R_{k+1}(2g(x_0)))$ -large set.

Consider the coloring $h : X \setminus \{x_0\} \rightarrow k+1$ defined by $h(y) = f(x_0, y)$. By Proposition 3.12(1), there exists some $(\omega^k \cdot R_{k+1}(2g(x_0)))$ -large and h -homogeneous subset Y of $X \setminus \{x_0\}$, and let $c_0 \leq k$ be the corresponding color.

By Lemma 3.13, as $\min Y \geq (k+1)^{R_{k+1}(2g(x_0))}$, there exists a sequence $Y_0 < \dots < Y_{g(x_0)-1}$ of $\omega^k + 1$ -large subsets of Y and some color c_1 such that $f(x, y) = c_1$ for every $x \in Y_s$ and $y \in Y_t$ for $s < t < g(x_0)$.

If $c_0 = c_1$, then the set $\{x_0\} \cup \{\min Y_0 < \dots < \min Y_{g(x_0)-1}\}$ is f -homogeneous and ω - g -large and we are done. So assume $c_0 \neq c_1$ and consider the coloring $f' : [X]^2 \rightarrow k$ to be the same coloring as f , except that the colors c_0 and c_1 are fused into one. By the inductive hypothesis, there exists some f' -homogeneous subsets $Z_1 \subseteq Y_1$.

If Z_1 is f homogeneous, then we are done. Otherwise, $f([Z_1]^2) = \{c_0, c_1\}$, and, since $\min Z_1 > R_2(g(\min Z_0))$ by sparsity of X , there exists some f -homogeneous subset $W \subseteq Z_1$ of size $\geq g(\min Z_0)$. If $f([W]^2) = \{c_0\}$, then $\{x_0\} \cup W$ is f -homogeneous, and if $f([W]^2) = \{c_1\}$, then $\{\min Z_0\} \cup W$ is f -homogeneous, and both of these sets are ω - g -large. \square

By generalizing the proof of Patey and Yokoyama [22, Lemma 4.4] to trRT_k^2 and replacing the bound of Ketonen and Solovay by Proposition 3.16, we obtain the following proposition:

Proposition 3.17. *Let $n \in \mathbb{N}$, $k \geq 1$ and let $X \subseteq \mathbb{N}$ be $(\omega^{kn} + 1)$ -large and $(x \mapsto (kn)^{R_{kn}(2x+2)})$ -sparse, then it is trRT_k^2 - ω^n -large.*

Proof. Consider a transitive coloring $f : [X]^2 \rightarrow k$ and let $\bar{f} : [X]^2 \rightarrow kn$ be defined by $\bar{f}(x, y) = kj + i$ if $f(x, y) = i$ and if j is the smallest index such that there is no ω^{j+1} -large subset $H \subseteq [x, y] \cap X$ with $x \in H$ and such that $H \cup \{y\}$ is f -homogeneous for color i , or, if no such index exists, take $j = n - 1$.

By Proposition 3.16, take $Y \subseteq X$ such that Y is ω - $(x \mapsto x+1)$ -large and \bar{f} -homogeneous. Write $Y = \{y_0 < y_1 < \dots < y_\ell\}$ for some $\ell > y_0$ and let $\bar{f}([Y]^2) = kj + i$. For every $s < \ell$, let $H_s \subseteq [y_s, y_{s+1})$ be ω^j -large, f -homogeneous for the color i and such that $y_s \in H_s$. Let $H = \{y_0\} \cup \bigcup_{s=1}^{\ell-1} H_s$, then, as $\ell > y_0$, H is ω^{j+1} -large and $H \cup \{y_\ell\}$ is f -homogeneous for the color i . Thus, we have $j = n$ as $\bar{f}(y_0, y_\ell) = nj + i$, hence H is f -homogeneous and ω^n -large. \square

Corollary 3.18. *Let $n \in \mathbb{N}$, $k \geq 1$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{nk+3} -large and $\min X \geq 8$, then it is trRT_k^2 - ω^n -large.*

Proof. Since $\min X \geq 8$, then by Lemma 3.8, there is an $\omega^2 \cdot 3$ -sparse $\omega^{nk} \cdot 2 + 1$ -large subset $Y \subseteq X$. By regularity, there are two subsets $Y_0 < Y_1$ of Y such that Y_0 is ω^{nk} -large and Y_1 is $\omega^{nk} + 1$ -large.

By Lemma 3.7, since Y_1 is $\omega^2 \cdot 3$ -sparse and $\min X \geq 7$, then it is $(x \mapsto x^{x^{x^{(2x+2)}}})$ -sparse. Since Y_0 is ω^{nk} -large, then $\text{card } Y_0 \geq nk$, so $\min Y_1 \geq nk$. It follows that Y_1 is $(x \mapsto (kn)^{(kn)^{kn(2x+2)}})$ -sparse, so by Proposition 3.17, Y_1 is trRT_k^2 - ω^n -large. \square

Corollary 3.19. *Let $n \in \mathbb{N}$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{2n+3} -large and $\min X \geq 9$, then it is ADS - ω^n -large.*

Proof. Immediate by Corollary 3.18. \square

Note that this bound is optimal, up to an additive constant, in that the coloring witnessing the lower bound of RT_2^2 by Kotlarski et al. [16, Theorem 5.4] is transitive.

3.8 Ramsey's theorem for pairs

We now have all the necessary ingredients to prove our optimal bounds for $\text{RT}_k^2\text{-}\omega^n$ -largeness, using the bounds for the fallow Erdős-Moser theorem and for transitive Ramsey's theorem for pairs. We first state it with largeness and sparsity assumptions, and then use Section 3.2 to compute purely largeness bounds.

Theorem 3.20. *Let $n, k \geq 1$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is $\omega^{kn} \cdot 4$ -large and $(x \mapsto x^{R_x(2x+2)})$ -sparse with $kn \leq \min X$, then it is $\text{RT}_k^2\text{-}\omega^n$ -large.*

Proof. Let $f : [X]^2 \rightarrow k$ be a coloring. X is $\omega^{kn} \cdot 4$ -large and $(x \mapsto x^{R_x(2x)})$ -sparse, hence, by Proposition 3.14(3), there exists some $\omega^{kn} + 1$ -large and f -fallow subset $Y \subseteq X \setminus \{\min X\}$. In particular, Y is f -transitive.

As $kn \leq \min X$, Y is $(x \mapsto (kn)^{R_{kn}(2x+2)})$ -sparse, and by Proposition 3.17 there exists some ω^n -large f -homogeneous subset $Z \subseteq Y$. \square

Note that if we applied fEM and trRT_k^2 under their explicit largeness assumptions (Corollaries 3.15 and 3.18) instead of their hybrid sparsity/largeness versions (Propositions 3.14 and 3.17), this would have yielded significantly worse bounds. It is therefore useful to keep a sparsity version of the statements in case of their use as an intermediate object in the proof of a stronger combinatorial statement. When translating the sparsity assumptions into largeness ones, we obtain our main theorem.

Main Theorem 1.2. *Let $n, k \geq 1$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{kn+3} -large and $\min X \geq 17$, then it is $\text{RT}_k^2\text{-}\omega^n$ -large.*

Proof. Since $\min X \geq 17$, then by Lemma 3.8, there is an $\omega^2 \cdot 3$ -sparse $\omega^{nk} \cdot 5 + 1$ -large subset $Y \subseteq X$. In particular, there are two subsets $X_0 < X_1$ of X such that X_0 is ω^{nk} -large and X_1 is $\omega^{nk} \cdot 4$ -large. Since X_0 is ω^{nk} -large, then $\text{card } X_0 \geq nk$, so $\min X_1 \geq nk$. Since X_1 is $\omega^2 \cdot 3$ -large, then by Lemma 3.7, it is $(x \mapsto x^{R_x(2x+2)})$ -sparse. It follows by Theorem 3.20 that X_1 is $\text{RT}_k^2\text{-}\omega^n$ -large. \square

References

- [1] Teresa Bigorajska and Henryk Kotlarski. A partition theorem for α -large sets. *Fund. Math.*, 160(1):27–37, 1999.
- [2] Teresa Bigorajska and Henryk Kotlarski. Some combinatorics involving ξ -large sets. *Fund. Math.*, 175(2):119–125, 2002.
- [3] Teresa Bigorajska and Henryk Kotlarski. Partitioning α -large sets: some lower bounds. *Trans. Amer. Math. Soc.*, 358(11):4981–5001, 2006.
- [4] Andrey Bovykin and Andreas Weiermann. The strength of infinitary ramseyan principles can be accessed by their densities. *Annals of Pure and Applied Logic*, 168(9):1700–1709, 2017.

- [5] Michiel De Smet and Andreas Weiermann. Partitioning α -large sets for $\alpha < \varepsilon_\omega$. *arXiv preprint arXiv:1001.2437*, 2010.
- [6] Damir D. Dzhafarov and Carl Mummert. *Reverse mathematics—problems, reductions, and proofs*. Theory and Applications of Computability. Springer, Cham, [2022] ©2022.
- [7] P. Erdős and L. Moser. On the representation of directed graphs as unions of orderings. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 9:125–132, 1964.
- [8] P. Erdős and G. Szekeres. A combinatorial problem in geometry. *Compositio Math.*, 2:463–470, 1935.
- [9] Ronald L. Graham, Bruce L. Rothschild, and Joel H. Spencer. *Ramsey theory*. Wiley Series in Discrete Mathematics and Optimization. John Wiley & Sons, Inc., Hoboken, NJ, 2013. Paperback edition of the second (1990) edition [MR1044995].
- [10] Godfrey H Hardy. A theorem concerning the infinite cardinal numbers. *Quarterly journal of Mathematics*, 35:87–94, 1904.
- [11] Denis R. Hirschfeldt and Richard A. Shore. Combinatorial principles weaker than Ramsey’s theorem for pairs. *J. Symbolic Logic*, 72(1):171–206, 2007.
- [12] Jussi Ketonen and Robert Solovay. Rapidly growing Ramsey functions. *Ann. of Math. (2)*, 113(2):267–314, 1981.
- [13] Leszek Aleksander Kołodziejczyk, Tin Lok Wong, and Keita Yokoyama. Ramsey’s theorem for pairs, collection, and proof size. *Journal of Mathematical Logic*, page 2350007, 2023.
- [14] Leszek Aleksander Kołodziejczyk and Keita Yokoyama. Some upper bounds on ordinal-valued Ramsey numbers for colourings of pairs. *Selecta Math. (N.S.)*, 26(4):Paper No. 56, 18, 2020.
- [15] Henryk Kotlarski. *A model-theoretic approach to proof theory*, volume 51 of *Trends in Logic—Studia Logica Library*. Springer, Cham, 2019. Edited by Zofia Adamowicz, Teresa Bigorajska and Konrad Zdanowski.
- [16] Henryk Kotlarski, Bożena Piekart, and Andreas Weiermann. More on lower bounds for partitioning α -large sets. *Ann. Pure Appl. Logic*, 147(3):113–126, 2007.
- [17] Quentin Le Houérou and Ludovic Levy Patey. Π_4^0 conservation of the Ordered Variable Word theorem. *The Journal of Symbolic Logic*, pages 1–16, 2023.
- [18] Quentin Le Houérou, Ludovic Levy Patey, and Keita Yokoyama. π_4^0 conservation of ramsey’s theorem for pairs. *Journal of the London Mathematical Society*, 113(1):e70419, 2026.

- [19] Manuel Lerman and Joseph G. Rosenstein. Recursive linear orderings. In *Patras Logic Symposium (Patras, 1980)*, volume 109 of *Stud. Logic Found. Math.*, pages 123–136. North-Holland, Amsterdam-New York, 1982.
- [20] Manuel Lerman, Reed Solomon, and Henry Towsner. Separating principles below Ramsey’s theorem for pairs. *J. Math. Log.*, 13(2):1350007, 44, 2013.
- [21] Ludovic Patey. *The reverse mathematics of Ramsey-type theorems*. PhD thesis, Université Paris Diderot (Paris 7) Sorbonne Paris Cité, 2016.
- [22] Ludovic Patey and Keita Yokoyama. The proof-theoretic strength of Ramsey’s theorem for pairs and two colors. *Adv. Math.*, 330:1034–1070, 2018.
- [23] Florian Pelupessy. On α -largeness and the paris-harrington principle in RCA_0 and RCA_0^* . 2016.
- [24] Z. Ratajczyk. A combinatorial analysis of functions provably recursive in $I\Sigma_n$. *Fund. Math.*, 130(3):191–213, 1988.
- [25] Stephen G. Simpson. *Subsystems of second order arithmetic*. Perspectives in Logic. Cambridge University Press, Cambridge; Association for Symbolic Logic, Poughkeepsie, NY, second edition, 2009.
- [26] Henry Towsner and Keita Yokoyama. Erdős-moser and $I\Sigma_2$. *Israel J. Math.*, 263(2):843–870, 2024.
- [27] S. S. Wainer. A classification of the ordinal recursive functions. *Arch. Math. Logik Grundlag.*, 13:136–153, 1970.